

New members in the Upper Sco association from the UKIDSS Early Data Release ^{*} †

N. Lodieu^{1‡}, N. C. Hambly², & R. F. Jameson¹

¹*Department of Physics & Astronomy, University of Leicester, University Road, Leicester LE1 7RH, UK*

²*Scottish Universities' Physics Alliance (SUPA), Institute for Astronomy, School of Physics, University of Edinburgh, Royal Observatory, Blackford Hill, Edinburgh EH9 3HJ, UK*

Accepted —. Received —; in original form —

ABSTRACT

We present the results of a 9.3 square degree infrared (*ZYJHK*) survey in the Upper Scorpius association extracted from the UKIRT Infrared Deep Sky Survey (UKIDSS) Galactic Cluster Survey Early Data Release. We have selected a total of 112 candidates from the (*Z*−*J*, *Z*) colour-magnitude diagram over the *Z* = 12.5–20.5 magnitude range, corresponding to *M* = 0.25–0.01 *M*_⊙ at an age of 5 Myr and a distance of 145 pc. Additional photometry in *J* and *K* filters revealed most of them as reddened stars, leaving 32 possible members. Among them, 15 have proper motion consistent with higher mass members from Hipparcos and optical spectra with strong H α in emission and weak gravity features. We have also extracted two lower mass candidate members for which no optical spectra are in hand. Three members exhibit strong H α equivalent widths (>20Å), suggesting that they could still undergo accretion whereas two other dwarfs show signs of chromospheric activity. The likelihood of the binarity of a couple of new stellar and substellar members is discussed as well.

Key words: Stars: low-mass, brown dwarfs — Techniques: photometric — Techniques: spectroscopic — Infrared: stars — open clusters and associations: individual: Upper Sco

1 INTRODUCTION

The Scorpius Centaurus is the nearest OB association, located at a mean distance of 145±2 pc from the Sun (de Bruijne et al. 1997). The association covers 120 square degrees and is composed of 3 subgroups: Upper Scorpius, Upper Centaurus Lupus, and Lower Centaurus Crux (Blaauw 1964; de Zeeuw et al. 1999). The age of the Upper Sco association is about 5 Myr with little scatter (Preibisch & Zinnecker 2002). The region is free of extinction with *A*_v ≤ 2 mag, suggesting that star formation has already ended (Walter et al. 1994).

The first survey conducted in Upper Sco was performed with the Einstein X-ray satellite by Walter et al. (1994), yielding 28 low-mass pre-main-sequence stars in 7 square degrees. Later on, de Bruijne et al. (1997) extracted 115 mem-

bers using kinematic information and trigonometric parallaxes for 1215 Hipparcos stars. A larger scale X-ray survey carried out by Preibisch et al. (1998) revealed several hundreds of X-rays sources confirmed spectroscopically as young pre-main-sequence members of the Scorpius Centaurus association. Deeper optical surveys complemented by near-infrared photometry (Ardila et al. 2000; Martín et al. 2004; Slesnick et al. 2006) crossed the hydrogen-burning limit with the discovery of members with spectral types later than M6 (Martín et al. 1996; Luhman et al. 1998). The current census of spectroscopic brown dwarfs in Upper Sco includes 46 M6, 10 M7, 6 M8, and 2 M9 dwarfs, all confirmed as members on the basis of their chromospheric activity and weak gravity features (Martín et al. 2004; Slesnick et al. 2006). Recently, we extended the cluster sequence down to 0.008 *M*_⊙ using UKIDSS Galactic Cluster Survey science verification data. A dozen new brown dwarf candidates with spectral types later than M9 and masses below 0.02 *M*_⊙ were extracted over 6.5 square degrees (Lodieu et al., *subm. to MNRAS*).

The UKIRT Infrared Deep Sky Survey (UKIDSS; Lawrence et al. 2006) constitutes the new generation of deep large-scale infrared surveys. The project is subdivided into 5

^{*} Based on observations made with the United Kingdom Infrared Telescope, operated by the Joint Astronomy Centre on behalf of the U.K. Particle Physics and Astronomy Research Council.

† Based on observations collected with the ESO 3.6-m/EFOSC2 at the European Southern Observatory, La Silla, Chile (ESO programme 076C-0237)

‡ E-mail: nl41@star.le.ac.uk




Figure 1. The UKIDSS GCS coverage in the Upper Sco association from the Early Data Release is depicted with boxes. Early-type stars from Hipparcos are displayed as 5-pointed star symbols (de Zeeuw et al. 1999). We have included known members: triangles are from Preibisch & Zinnecker (2002), diamonds from Ardila et al. (2000), squares from Martín et al. (2004), and pluses from Slesnick et al. (2006). The new spectroscopic members and non-members from this study are shown as filled circles and star symbols, respectively. Diamonds located in the area surveyed by the EDR are potential reddened members. Filled circles outside the boxes represent new members of the association extracted from the UKIDSS GCS science verification (Lodieu et al. 2006). Spectroscopic non-members are located in an area affected by reddening at $(RA, dec) \sim (250^\circ, -21.8^\circ)$.

Table 1. Log of the observations. Central coordinates (in J2000) of each WFCAM tile is provided along with the observing date. Each tile was observed in all five passbands (ZYJHK).

Tile	R.A.	Dec.	Date
1	15 42	-21 36	2005-06-05
2	15 46	-21 38	2005-06-05
3	15 50	-21 38	2005-06-05
4	15 53	-21 38	2005-06-05
5	16 13	-25 16	2005-06-11
6	16 25	-25 03	2005-06-14
7	16 28	-21 36	2005-06-05
8	16 31	-21 38	2005-06-05
9	16 35	-21 38	2005-06-05
10	16 36	-25 16	2005-06-06
11	16 39	-21 38	2005-06-05
12	16 40	-25 09	2005-06-06

surveys: The Large Area Survey, the Galactic Cluster Survey (GCS), the Galactic Plane Survey, the Deep Extragalactic Survey, and the Ultra-Deep Survey. The GCS will survey over thousand square degrees in 10 star-forming regions and open clusters down to $K = 18.4$ mag at two epochs. The

main goal of the GCS is to study the Initial Mass Function well down into the substellar regime in an homogeneous manner in 10 star-forming regions and open clusters.

In this paper we concentrate on the selection of new cluster member candidates in a 9.3 square degree area in the Upper Scorpius association from the UKIDSS Early Data Release (EDR; Dye et al. 2006). The photometric, proper motion, and spectroscopic results presented in this paper represent the first step towards the ultimate goal of the GCS. In Sect. 2 we briefly present the infrared photometric observations conducted in Upper Sco. In Sect. 3 we describe the selection of new candidates from various colour-magnitude diagrams complemented by proper motion. In Sect. 4 we present the optical spectroscopic follow-up carried out with the ESO 3.6-m telescope. We infer spectral types and measure equivalent widths for the $H\alpha$ emission line and NaI doublet to assess the membership of the new candidates. Finally, we summarise our work in Sect. 5.

2 THE UKIDSS GCS IN UPPER SCO

2.1 The Early Data Release (EDR)

Twelve WFCAM tiles were observed in the association during the first phase of survey observations in June 2005 (Table 1), covering a total of 9.3 square degrees (Fig. 1) in *ZYJHK* broad-band filters (Hewett et al. 2006). A summary of the observations, including the 12 WFCAM tiles, central coordinates and date of observations is provided in Table 1. Uniformity of quality in the UKIDSS EDR (e.g. seeing, photometricity, etc.) is discussed at length in Dye et al. (2006). We refer the reader to Lawrence et al. (2006) for more details on the observing strategy of the UKIDSS¹ project and its individual components.

All observations carried out within the framework of the UKIDSS project are pipeline-processed at the Cambridge Astronomical Survey Unit (CASU; Irwin et al. 2006, in prep.)². The processed data are then archived in Edinburgh and released to the European community through the WFCAM Science Archive (WSA; Hambly et al. 2006, in prep.)³.

2.2 Observations and colour-magnitude diagrams

The UKIDSS GCS observations taken in Upper Sco and included in the Early Data Release amounts for a total of 9.3 square degrees (Fig. 1). There are 3 main blocks centered approximately at (RA,dec)=(237.0,−21.55), (248.0,−21.55), and (250.05,−25.15) with three additional small blocks (made of 2 pawprints) contributing for 0.3 deg² in total. The survey was conducted in *ZYJHK* and reached 5 σ completeness limits of $Z = 20.0$, $Y = 19.6$, $J = 18.6$, $H = 18.1$, and $K = 17.5$ mag. Hence, this survey is 3 magnitudes deeper than any previous study in the region and is able to probe young L dwarfs according to the DUSTY 5 Myr isochrones. The two lowest mass spectroscopic members reported in Upper Sco to date have spectral types of M9 corresponding to ~ 20 Jupiter masses (Martín et al. 2004; Slesnick et al. 2006).

We have selected from the EDR point sources (Class parameter set to -1 or -2 in all passbands) detected in all 5 filters (*ZYJHK*) by use of the SQL script given in Appendix A injected in the WFCAM Science Archive. The total number of point sources is 174,010 (dots in the colour-magnitude diagrams in Fig. 2) in the $Z = 11.4$ – 20.5 magnitude range, corresponding to masses between $0.35 M_{\odot}$ and $0.008 M_{\odot}$ at an age of 5 Myr and a distance of 145 pc (Baraffe et al. 1998; Chabrier et al. 2000).

The colour-magnitude diagrams resulting from the GCS observations in Upper Sco are shown in Fig. 2, including the $(Z - J, Z)$, $(Z - K, Z)$, and $(J - K, J)$ diagrams. Overplotted are 5 and 10 Myr NextGen (solid lines; Baraffe et al. 1998), DUSTY (dashed lines; Chabrier et al. 2000), and COND (dotted lines; Baraffe et al. 2002) isochrones shifted at the distance of the association ($d = 145$ pc). Note that

isochrones were computed for the UKIDSS passbands (courtesy I. Baraffe and F. Allard).

3 NEW CANDIDATE MEMBERS OF THE UPPER SCO ASSOCIATION

This section describes the selection of new candidates in the Upper Scorpius association using colour-magnitude diagrams and proper motions.

3.1 Selection of candidates in Upper Sco

Candidates in open clusters are generally selected to the right of the theoretical isochrones or the Zero-Age-Main-Sequence (Leggett 1992). To select all possible members in Upper Sco at the expense of including some contaminants, we have extracted all candidates to the right of a straight line running from $(Z - J, Z) = (0.8, 11.5)$ to $(2.1, 21.5)$. The total number of candidates is 192. However, we have limited our analysis to objects fainter than $Z = 12.5$ ($J \sim 11.5$), leaving 112 candidates. This threshold translates into a mass of $0.25 M_{\odot}$ and corresponds to completeness limit of the survey conducted by Preibisch & Zinnecker (2002) in the region. We provide an electronic table with all photometric candidates, including those classified later as non-members after proper motion and spectroscopic analysis (Table 2). Some candidates fainter than $Z = 17$ mag are running along the straight line defined as our criterion and located to the left of the theoretical isochrones (Chabrier et al. 2000) in the $(Z - J, Z)$ diagram. Consequently, we cast doubt on the membership of those objects and do not consider them in the subsequent analysis for two reasons: first, one candidate located at $Z \sim 16.2$ and $Z - J \sim 1.5$ turned out as a non-member after optical spectroscopy, and, second, the cluster sequence runs further to the red and to the right of the isochrones as demonstrated in Lodieu et al. (2006, subm. to MNRAS).

To further assess the membership of the 112 candidates, we have investigated their location in the $(Z - K, Z)$ and $(J - K, J)$ colour-magnitude diagrams (top right and bottom left graphs in Fig. 2, respectively) as well as in the $(Z - K, J - K)$ two-colour diagram. The $(Z - K, Z)$ diagram is characterised by a spread of the sample of candidates which can be explained by the presence of interstellar reddening in the part of Upper Sco released in the Early Data Release. This trend is confirmed in the $(J - K, J)$ diagram where we indicated a reddening vector of $A_V = 1$ and by the location of those sources in Fig. 1. As a consequence, we have applied the following colour criteria to weed out reddened sources:

- $0.90 \leq J - K \leq 1.10$ for $J = 11.5$ – $13.0 \Rightarrow 23$ candidates
- $0.90 \leq J - K \leq 1.25$ for $J = 13.0$ – $14.0 \Rightarrow 5$ candidates
- All 4 sources fainter than $J = 14$ mag were kept

A total of 32 sources remain as photometric candidates. Their membership is further investigated using proper motions (Sect. 3.2) and optical spectroscopy (Sect. 4) as additional criteria.

¹ The main page is www.ukidss.org

² The CASU WFCAM webpage can be found at <http://apm15.ast.cam.ac.uk/wfcam>

³ The WFCAM Science Archive is accessible at <http://surveys.roe.ac.uk/wsa>

Table 2. All 112 photometric candidates selected from the $(Z - J, Z)$ colour-magnitude diagram in 9.3 square degrees in the Upper Sco association. This table provides coordinates (J2000), $ZYJHK$ photometry from the GCS, and proper motion from the GCS/2MASS cross-correlation for members and non-members. The table is available in electronic format only.

R.A.	Dec	Z	Y	J	H	K	$\mu_\alpha \cos \delta$	μ_δ
16:26:38.63	-21:58:51.7	12.803	12.246	11.721	11.061	10.696	-18.32	-21.49
...	...	...	...	...	...	...	...	...
...	...	...	...	...	...	...	...	...
16:28:52.80	-21:47:41.1	12.819	12.266	11.799	11.109	10.802	-26.03	-7.87

Table 3. Seventeen new members of the Upper Sco association confirmed from their photometry, proper motion, and spectroscopy (filled circles in Fig. 2). This table lists the name of the object according to the IAU nomenclature, the equatorial coordinates (J2000), the magnitudes from the GCS, the proper motions (in mas/yr), the $H\alpha$ and NaI equivalent widths (in Å), and the spectral types with an accuracy of half a subclass. The two faintest candidates ($Z > 17.0$ mag) have neither spectral types nor equivalent width measurements because they are too faint for optical spectroscopy with ESO 3.6-m/EFOSC2 but one has a proper motion consistent with the association.

USco J...	R.A.	Dec	Z	Y	J	H	K	$\mu_\alpha \cos \delta$	μ_δ	$H\alpha$	NaI	SpT
1626-2158	16:26:38.63	-21:58:51.7	12.803	12.246	11.721	11.061	10.696	-18.32	-21.49	-4.70	2.56	M4.0
1628-2147	16:28:52.80	-21:47:41.1	12.819	12.266	11.799	11.109	10.802	-26.03	-7.87	-4.40	3.55	M3.5
1549-2146	15:49:22.59	-21:46:57.6	13.074	12.536	12.019	11.387	11.116	-26.11	-15.32	-9.10	4.40	M3.5
1553-2114	15:53:01.32	-21:14:13.7	13.081	12.486	11.927	11.329	10.983	-10.63	-25.37	-30.70	2.07	M4.0
1551-2146	15:51:40.32	-21:46:10.6	13.094	12.476	11.931	11.382	10.966	-7.39	-27.57	-9.00	2.41	M4.0
1551-2145	15:51:18.70	-21:45:23.7	13.288	12.651	12.102	11.472	11.177	-14.48	-18.23	-9.60	3.12	M4.0
1627-2138	16:27:25.52	-21:38:03.8	13.384	12.751	12.165	11.531	11.147	-11.60	-22.38	-5.90	3.38	M4.0
1629-2137	16:29:48.78	-21:37:08.9	13.606	12.957	12.407	11.845	11.444	-18.79	-33.66	-30.00	3.25	M5.0
1555-2117	15:55:05.31	-21:17:40.4	13.815	13.136	12.503	11.900	11.534	-14.15	-18.44	-13.00	2.42	M5.0
1547-2137	15:47:04.94	-21:37:40.5	13.885	13.230	12.697	12.051	11.728	2.06	-18.06	-10.80	3.78	M4.0
1634-2201	16:34:28.50	-22:01:12.1	14.760	13.953	13.260	12.612	12.108	-8.80	-23.06	-13.60	3.70	M6.0
1549-2201	15:49:57.33	-22:01:25.7	14.867	13.989	13.361	12.792	12.351	-6.68	-13.67	-12.30	2.88	M6.0
1613-2509	16:13:48.80	-25:09:00.9	14.974	14.202	13.548	12.999	12.470	-6.98	-29.66	-25.50	2.37	M5.0
1549-2120	15:49:04.14	-21:20:15.2	15.204	14.432	13.770	13.212	12.827	-11.28	-15.79	-23.00	3.20	M6.0
1554-2135	15:54:19.99	-21:35:43.1	16.737	15.726	14.932	14.282	13.708	-5.58	-27.06	-29.40	2.80	M8.0
1547-2139	15:47:22.82	-21:39:14.3	17.940	16.559	15.637	14.834	14.177	-6.74	-15.99	—	—	—
1639-2534	16:39:19.15	-25:34:09.9	19.942	18.328	17.202	16.395	15.607	—	—	—	—	—

Table 4. This table lists the equatorial coordinates (in J2000), the $ZYJHK$ magnitudes, and the proper motion of 15 photometric candidates classified as non-members after optical spectroscopic follow-up (star symbols and open triangles in Fig. 2). The spectra look like reddened early-type stars apart from two objects classified as a M5 dwarf (15 50 39.55, -21 39 47.5) and a late-M dwarf (15 50 11.5, -22 01 21.9). All spectroscopic non members but 3 exhibit proper motions discrepant with the mean motion of the association.

R.A.	Dec.	Z	Y	J	H	K	$\mu_\alpha \cos \delta$	μ_δ
16:37:38.13	-21:52:50.2	12.516	12.097	11.566	10.838	10.511	-10.17	-5.66
16:37:52.30	-22:01:43.9	12.530	12.054	11.506	10.793	10.429	-3.03	-7.79
16:35:21.87	-22:03:12.0	12.609	12.139	11.601	10.798	10.503	-7.20	-0.09
16:39:45.13	-21:36:19.0	12.667	12.202	11.683	11.004	10.682	21.95	-4.01
16:39:06.86	-22:01:33.8	12.723	12.233	11.710	10.967	10.668	-0.81	-5.75
16:34:56.41	-21:46:04.0	12.731	12.255	11.704	10.926	10.633	-1.98	-6.97
16:35:25.71	-21:53:15.4	12.735	12.253	11.755	11.041	10.662	-3.11	-5.48
16:39:40.12	-21:38:42.6	12.775	12.304	11.783	11.025	10.742	15.98	-6.72
16:36:33.90	-22:01:08.4	12.979	12.453	11.915	11.237	10.906	-7.13	-2.83
16:36:40.01	-22:01:46.4	13.145	12.571	11.998	11.291	10.961	7.36	-2.03
15:50:39.55	-21:39:47.5	13.245	12.587	12.071	11.465	11.137	-16.16	-21.77
16:32:59.62	-21:51:41.9	13.396	12.877	12.320	11.511	11.249	-13.57	0.85
16:36:51.29	-21:39:08.3	13.786	13.111	12.632	12.026	11.837	-1.56	-0.41
16:35:00.85	-21:16:55.3	15.876	14.708	13.853	13.273	12.651	-35.45	-48.37
15:50:11.50	-22:01:21.9	17.202	16.283	15.572	14.969	14.543	-20.47	-80.92




Figure 2. Colour-magnitude diagrams for 9.3 square degrees in the Upper Scorpius association from the UKIDSS Early Data Release. Large filled circles are new spectroscopic Upper Sco members. Open triangles and star symbols are spectroscopic and proper motion non-members, respectively. The dots represent all candidates classified photometrically as reddened stars from their location in the $(J - K, J)$ and $(Z - K, J - K)$ diagrams. The diamonds are possible reddened members for which no optical spectroscopy is currently available. Overplotted are the 5 and 10 Myr NextGen (solid line; Baraffe et al. 1998), DUSTY (dashed line; Chabrier et al. 2000), and COND (dotted line; Baraffe et al. 2002) isochrones. The mass scale is shown on the right hand side of the diagrams and spans $0.3\text{--}0.008 M_{\odot}$. *Top left:* $(Z - J, Z)$ diagram. *Top right:* $(Z - K, Z)$ diagram. *Bottom left:* $(J - K, J)$ diagram. *Bottom right:* Vector point diagram for sources brighter than $J = 15.8$ mag because the faintest ones do not have proper motion measurements due to the lack of 2MASS detection. The last three diagrams contain only photometric candidates selected in the $(Z - J, Z)$ diagram.

3.2 Proper motions

Optical or near-infrared photometry alone is not sufficient to extract a clean sample of members in open clusters. We have exploited the 2MASS database and the GCS observations as first and second epoch, respectively, to compute

proper motions for candidates brighter than $J = 15.8$ mag, corresponding to the 2MASS 5σ completeness limit and a mass of $0.015 M_{\odot}$ in Upper Sco (Chabrier et al. 2000). The vector point diagram (proper motion in right ascension versus proper motion in declination) is presented in the bottom right panel in Fig. 2. Most of the sources classified as

Table 5. This table lists the equatorial coordinates (in J2000), the *ZYJHK* magnitudes, and the proper motion of 15 potential reddened members in Upper Sco whose proper motion is consistent with the association (diamonds in Fig. 2).

R.A.	Dec.	<i>Z</i>	<i>Y</i>	<i>J</i>	<i>H</i>	<i>K</i>	$\mu_\alpha \cos \delta$	μ_δ
16:38:26.94	-22:03:55.8	12.541	11.997	11.419	10.680	10.248	-1.687	-13.701
16:13:36.51	-25:03:47.4	12.563	12.056	11.456	10.614	10.262	+5.603	-15.645
16:26:39.26	-21:13:45.4	13.133	12.451	11.859	11.133	10.704	-14.451	-8.276
16:37:50.68	-22:03:07.9	13.143	12.612	12.105	11.292	10.963	-7.985	-9.292
16:38:01.92	-22:01:18.9	13.482	12.970	12.389	11.461	11.138	+0.675	-9.565
16:36:56.95	-22:02:32.1	13.550	13.026	12.439	11.541	11.204	-14.490	-7.445
16:35:53.43	-21:56:31.2	14.052	13.138	12.155	11.046	10.387	-2.271	-8.968
16:36:32.01	-22:03:13.0	14.108	13.070	11.996	10.644	9.965	-14.992	-5.803
16:36:58.40	-22:04:12.1	14.113	13.505	12.898	11.912	11.605	-20.965	-8.786
16:35:37.41	-21:49:36.3	14.123	13.519	12.914	12.019	11.659	-12.573	-12.365
16:27:09.41	-21:48:45.7	14.316	13.529	12.876	12.113	11.659	-13.443	-7.843
15:51:47.09	-21:13:23.6	15.000	14.310	13.543	12.440	11.516	-12.856	-23.383
16:35:59.71	-21:45:59.0	15.194	14.441	13.670	12.484	12.061	+0.691	-28.230
16:38:48.92	-22:01:07.0	15.506	14.793	14.056	12.965	12.572	-14.298	-10.709
15:52:10.88	-21:25:37.5	15.611	14.729	13.840	13.068	12.041	-9.209	-36.159

reddened stars exhibit a small proper motion (small dots), hence confirming their status of distant objects and non-members. However, there remain 15 objects (diamonds; Table 5) considered as reddened stars in the previous section which do have a proper motion within the 2σ circle plotted in the vector point diagram in Fig. 2. Although most of them are likely to be reddened stars because they lie in the same part of the diagram as the spectroscopic members (Fig. 1), they might be some reddened members. Unfortunately we do not provide optical spectroscopy for them but they need to be followed-up to determine how complete is the sample selected from the sequence alone. Two other groups of sources for which we obtained optical spectra (see Sect. 4 for more details) are overplotted. On the one hand, 12 candidates classified as non-members and plotted as star symbols in Fig. 2 have small proper motion close to $(\mu_\alpha \cos \delta, \mu_\delta) = (0,0)$ mas/yr. On the other hand, 20 new candidates shown as filled circles and triangles in Fig. 2 exhibit proper motion consistent with the mean motion of the association from Hipparcos (de Bruijne et al. 1997; Preibisch et al. 1998). The faintest photometric member candidate is too faint to compute its proper motion from the 2MASS images.

4 SPECTROSCOPIC FOLLOW-UP

This section describes the spectroscopic observations of new candidates in Upper Sco and discuss their membership.

4.1 Spectroscopic observations

We have carried out low-resolution optical spectroscopy of 31 candidates in Upper Sco with EFOSC2 on the ESO 3.6-m telescope in La Silla, Chile. The observations were obtained over 5 nights on 27 March-01 April 2006. Conditions were photometric during the first three nights whereas the last two nights were affected by thin cirrus. Seeing was typical in the 0.5–1.0 arcsec range during the observing run.

The EFOSC2 detector is a 2048×2048 pixel MIT CCDs with a pixel size of $0.157''/\text{pix}$, yielding a 5.5×5.5 arcmin field-of-view. We have employed the grism #16

with a 1 arcsec slit to cover the wavelength range from 6000 to 10300 Å at a resolution of $R \sim 300$. We have obtained an internal flat-field after each spectrum to remove fringing longwards of 7500 Å. Each target was observed once with exposure times ranging from 240 sec for the brightest objects to 3600 sec for the faintest objects, respectively. In addition, we have observed 3 known Upper Sco members: SCH160147–244101 (M5; Slesnick et al. 2006), DENIS J160514–240653 (M6; Martín et al. 2004), and SCH162528–165851 (M8; Slesnick et al. 2006) as well as a M5 field dwarf (SCH160311–134544) classified as a non-member by Slesnick et al. (2006). A couple of other field dwarfs were observed, including Gl643 (M3.5), Gl699 (M4), Gl166 (M4.5), GJ406 (M6), and vB10 (M8). Spectrophotometric standard stars (HR4963, LTT3218, LTT3864, and EG274; Hamuy et al. 1992) were observed during the night with slits of 1 and 5 arcsec to calibrate our targets. Bias, dome flats, and arc lamps were observed during the afternoon before each night.

4.2 Data reduction

The data reduction was carried out under the IRAF environment following standard procedures. First, we combined 10 bias frames and subtracted the averaged bias from the combined flat field and the science target. Second, we have divided the science frame by the normalised response function of the internal flat field observed immediately after each target. Afterwards, we extracted a one-dimensional spectrum interactively with apsum by choosing the appropriate aperture and background intervals. Then, we used the arc lamps to calibrate our spectra, yielding typical rms better than $0.35\text{\AA}$. The flux calibration with the spectrophotometric standard stars was not possible due to the presence of 2nd order contamination longwards of 7200 Å. All four standard targeted during the observing are white dwarfs or early-type stars, yielding an overestimate of the detector response in the red part of the spectrum. To solve this issue,

we used the optical spectra of 3 Upper Sco members⁴ to correct the shape of our spectra. However, the amount of flux from the blue end of the M dwarf spectra kicks in redwards of $\sim 9000\text{\AA}$, making the calibration very difficult. Thus, the shape of the science target and then comparison with templates is only valid between $6400\text{\AA}$ and $8500\text{\AA}$. No correction for telluric bands was made to the spectra. The optical spectra (normalised at $7500\text{\AA}$) of Upper Sco members are displayed in Fig. 3.

4.3 Spectral classification

The classification of M dwarfs is generally based on spectral indices measuring the strength of molecular absorption bands and atomic lines (Martín et al. 1999; Kirkpatrick et al. 1999). Spectral indices are defined over a large wavelength range and are valid for old field dwarfs. The effect of gravity can influence the computation of the indices (Martín et al. 1996). Thus, we preferred to rely on the direct comparison with known Upper Sco members and field M dwarfs observed with the same telescope/instrument set-up in order to classify our targets. We used 3 known Upper Sco members, SCH1601–2441 (M5; Slesnick et al. 2006), DENIS1605–2406 (M6; Martín et al. 2004), and SCH1625–1658 (M8; Slesnick et al. 2006), as young M dwarf templates.

The spectral classification of all candidates followed-up spectroscopically revealed 3 subgroups (we do not have spectra for the two faintest candidates):

(i) 12 proper motion non-members (star symbols in Fig. 2) classified photometrically as reddened sources are indeed spectroscopic non-members. All but two sources are reddened early-type stars. They are clearly concentrated in one part of the association (Fig. 1) associated with reddening and possibly due to the presence of a nearby early-type star. The presence of interstellar absorption is seen in the IRAS $100\mu\text{m}$ map of the association displayed in Fig. 1 in Slesnick et al. (2006). The other two sources are classified as a M5 dwarf ($15^{\text{h}}50^{\text{m}}39.55^{\text{s}}$, $-21^{\circ}39'47.5''$) and the remaining one is a late-M dwarf ($15^{\text{h}}50^{\text{m}}11.5^{\text{s}}$, $-22^{\circ}01'21.9''$). The latter is redder than the coolest Upper Sco member observed in this study (spectral type of M8) and exhibits $\text{H}\alpha$ in emission. The NaI equivalent width ($7.3\text{\AA}$) is larger than measurements for Upper Sco members and its proper motion is inconsistent with the association. It is also redder than vB10 (M8; Kirkpatrick et al. 1991) which we included in our program as a field M dwarf template. We tentatively classify this source as a M9 dwarf.

(ii) 3 proper motion members classified as spectroscopic non-members (triangles in Fig. 2): two turned out to be early-type stars, and the remaining one is a M5 dwarf.

(iii) 15 proper motion members confirmed spectroscopically as young M dwarfs belonging to the Upper Sco association. We can subdivide those members into five groups (Table 3; Fig. 3): 2 objects are classified as M3.5 dwarfs, 6 as M4.0, 5 as M5, 3 as M6, and the faintest object is a M8 dwarf. We assign an uncertainty of half a subclass to




Figure 4. $\text{H}\alpha$ equivalent widths as a function of spectral type for 15 new spectroscopic members (filled circles) and non-members (asterisks). The level of chromospheric activity confirms that those candidates as members of the Upper Sco association. We have also included equivalent widths for members (open triangles) published by Slesnick et al. (2006) as well as for field dwarfs observed with EFOSC2 and show $\text{H}\alpha$ emission (squares). Overplotted is the accretor/non-accretor empirical function (solid line) defined by Barrado y Navascués & Martín (2003). Three objects with spectral types of M4 and M5 lie above the solid line and could be accretors.

our spectral types (note that we rely on previous classifications). Young cluster members with spectral types of M6 or later are considered as brown dwarfs (Martín et al. 1996; Luhman et al. 1998). All but one spectroscopic members lie within a 2σ circle centred on $(-11, -25)$ mas/yr, assuming 10 mas/yr as typical errors on the proper motion measurement.

Consequently, 15 out of 18 proper motion members were confirmed as spectroscopic members, yielding a contamination of $3/18 = 16.5\%$ among our sample, lower than studies in the Pleiades (31%; Moraux et al. 2001) and α Per (30–45%; Barrado y Navascués et al. 2002) open clusters.

4.4 Chromospheric activity and gravity

To assess the youth of the M dwarfs having photometry and proper motion consistent with Upper Sco, we investigated the pseudo-equivalent widths of the $\text{H}\alpha$ emission line ($6365\text{\AA}$) and the NaI doublet ($8183/8195\text{\AA}$).

The $\text{H}\alpha$ equivalent widths measured for all Upper Sco members range from $-4.4\text{\AA}$ to $-30.7\text{\AA}$ (filled circles in Fig. 4) and lie below the empirical boundary between accreting and non accreting low-mass and brown dwarfs defined by Barrado y Navascués & Martín (2003). Three objects with strong $\text{H}\alpha$ emission lines lie above this line and are discussed in Sect. 4.5. Finally, we confirm the increasing strength of the $\text{H}\alpha$ equivalent width with later spectral type.

⁴ Spectra kindly provided by Eduardo Martín and Catherine Slesnick




Figure 3. Optical (6400–8500Å) spectra of 15 new members in the Upper Sco association obtained with EFOSC on the ESO 3.6-m telescope. The spectral types range from M3.5 to M8 with a typical uncertainty of half a subclass. We have added three known members for comparison purposes: SCH160147–244101 (M5; Slesnick et al. 2006), DENIS J160514–240653 (M6; Martín et al. 2004), and SCH162528–165851 (M8; Slesnick et al. 2006). All objects show strong H α emission and weak NaI doublets, indicator of low gravity.




Figure 5. NaI equivalent widths as a function of spectral type for 15 spectroscopic new members (filled circles) and non-members (asterisks). Three known members published by Martín et al. (2004) and Slesnick et al. (2006) were chosen as spectroscopic templates and are plotted as diamonds. Equivalent widths for 5 field dwarfs targeted with the same telescope and instrument configurations as the Upper Sco sample are displayed as squares. The weak NaI doublet equivalent widths measured for the Upper Sco candidate members indicate low gravity and adds credence to their membership.

The NaI doublet is a good gravity indicator for spectral types later than about M3 as it weakens with lower gravity i.e. younger ages. The equivalent widths measured for the Upper Sco members (filled circles) and field dwarfs (squares) observed with the same instrument are plotted as a function of spectral type in Fig. 5. Our measurements indicate that young Upper Sco members exhibit weaker NaI doublets than field dwarfs of similar spectral type. To put our work into a wider picture, our equivalent width measurements tend to be weaker than values reported in the literature for the Pleiades (Martín et al. 1996) although the spectral resolution was different. This argument remains however qualitative as lower resolution tend to overestimate equivalent widths.

To summarise, 15 out of 18 photometric and proper motion selected members have confirmatory signatures of weak NaI and H α emission (see optical spectra in Fig. 3). Our measurements are in agreement with results obtained for three known Upper Sco members published by Martín et al. (2004) and Slesnick et al. (2006) that we observed as templates during our programme.

4.5 Activity and accretion

Two different mechanisms can account for the H α emission line detected in all Upper Sco members (Fig. 4): chromospheric activity is characterised by weak lines with gaussian




Figure 6. Zoom on the NaI doublet at 8183/8195 Å for field dwarfs (black) and Upper Sco candidates (red) of same spectral type: M6 (bottom panel) and M4 (top panel). The NaI doublet is gravity sensitive i.e. the younger the object, the weaker the equivalent width is.

profiles whereas accretion and winds present as strong emission with asymmetric line shapes (Mohanty et al. 2005).

We have measured H α equivalent widths which are discrepant from values reported in the literature for spectroscopic templates. We attribute this variation to chromospheric activity. We should nevertheless emphasise that the difference in spectral resolution between our observations and those published in the literature could account for some of the discrepancy. The first object, DENIS J160514–240653 (Martín et al. 2004), was included in our program as a known Upper Sco member. We measured an equivalent width of -15.7Å in our spectrum compared to -24Å reported in Martín et al. (2004). The second object is SCH160311–13454, classified as a M5 field dwarf by Slesnick et al. (2006). We have measured an H α equivalent width of -18.8Å , stronger than the one reported previously (-8.7Å). This type of variation in the H α is common among mid-M field dwarfs (Hawley et al. 1996). We have also investigated the NaI doublet for this object and measured equivalent widths of 5.5Å and 7.0Å in our spectrum and Slesnick’s spectrum, respectively. Those values are suggestive of an age older than Upper Sco. We also measured a proper motion of $(+90, -54)$ mas/yr between the 2MASS observations and the optical survey by Slesnick et al. (2006), clearly inconsistent with the association.

The empirical relation defined by Barrado y Navascués & Martín (2003) from low-resolution optical spectroscopy to distinguish accreting and non-accreting low-mass stars and brown dwarfs is overplotted in Fig. 4. Most members lie below this empirical relation. However, three objects, one M4 and two M5 dwarf members of Upper Sco exhibit H α equivalent widths two to three times those of sources with the same spectral type and larger than the divide line between accretors and non-accretors. Although weaker than the strongest H α equivalent widths reported by Slesnick et al. (2006) in several objects (open triangles in Fig. 4) and despite our low spectral resolution which might have overestimated the equivalent widths, those

sources could still undergo accretion unless we observed strong flares due to chromospheric activity. Higher resolution spectroscopy centred on $H\alpha$ is required to investigate the line profiles and the origin of the strong emission.

4.6 Binarity

Bouy et al. (2006) reported a possible excess of wide binaries with separations between 100 and 150 AU among low-mass stars and brown dwarfs in Upper Sco. Similarly, Luhman (2005) discovered serendipitously a wide low-mass pair with a projected physical separation of 130 AU in the association.

A couple objects brighter than $J = 14$ mag lie clearly to the right of the cluster sequence which runs almost vertically in the $(J - K, J)$ colour-magnitude diagram, suggesting that they are possible low-mass binaries. The object at $J = 15.5$ mag ($M \sim 0.015 M_{\odot}$) could also be a multiple substellar system and of particular interest to test evolutionary models at low masses (if confirmed as such). Other binary candidates might be hidden in the list of potential reddened members. Because we selected only point sources in the EDR catalogue, we are not sensitive to separations of about 1-2 pixel or 0.4–0.8 arcsec. This corresponds to projected physical separations of 50–100 AU at the distance of the association ($d = 145$ pc). We are also insensitive to closer companions as it requires high-resolution imaging or adaptive optics although the location of objects in the various colour-magnitude diagrams can give some clues about the possible multiplicity of a source. Finally, we should be sensitive to larger separations and be able to detect wide companions on seeing-limited images (separation larger than a few hundreds of astronomical units) especially on the K -band images where microstepping is used to measure accurate proper motions when the second epoch observations are released.

5 SUMMARY

We have presented the analysis of a 9.3 square degree survey in the Upper Scorpius association conducted for the UKIDSS Galactic Cluster Survey and extracted from the Early Data Release. Combining infrared photometry, proper motion, and optical spectroscopy, the membership of the original 112 candidates can be summarised as follows:

(i) 80 photometric candidates selected from the $(Z - J, Z)$ colour-magnitude diagram were rejected on the basis their $J - K$ affected by reddening. Among them, 15 sources could be reddened members as they exhibit proper motion consistent with the association

(ii) 13 photometric candidates have proper motions inconsistent with the Upper Sco association and optical spectra of early-type stars affected by reddening

(iii) 3 objects have optical spectroscopy inconsistent with membership despite their photometry and proper motion suggesting that they are possible members

(iv) 15 candidates are definite members confirmed from photometry, proper motion, and optical spectroscopy, including 4 new brown dwarfs (spectral types later than M6). They span masses between 0.2 and $0.01 M_{\odot}$, assuming an age of 5 Myr and a distance of 145 pc for Upper Sco

(v) 2 faint photometric candidates with estimated

masses below $20 M_{\text{Jup}}$ according to the DUSTY models (Chabrier et al. 2000). One object has proper motion consistent with Upper Sco and no spectroscopy whereas the other one has neither proper motion nor spectroscopy currently available

We detected variation in the $H\alpha$ equivalent widths for one Upper Sco members and one field dwarf, hence showing signs of chromospheric activity. In addition, 3 new members exhibit $H\alpha$ emission lines stronger than the lower limit of the empirical relation between accreting and non accreting low-mass stars and brown dwarfs. Finally, a handful of members present a high probability of being binary systems from their location in the colour-magnitude diagrams.

ACKNOWLEDGMENTS

NL is a postdoctoral research associate funded by PPARC UK. We are grateful to Isabelle Baraffe and France Allard for providing us with the NextGen and DUSTY models for the WFCAM filters and Eduardo Martín and Catherine Slesnick for kindly supplying their optical spectra. We thank our colleagues at the UK Astronomy Technology Centre, the Joint Astronomy Centre in Hawaii, the Cambridge Astronomical Survey and Edinburgh Wide Field Astronomy Units for building and operating WFCAM and its associated data flow system. This research has made use of the Simbad database, operated at the Centre de Données Astronomiques de Strasbourg (CDS), and of NASA's Astrophysics Data System Bibliographic Services (ADS). This publication makes use of data products from the Two Micron All Sky Survey, which is a joint project of the University of Massachusetts and the Infrared Processing and Analysis Center/California Institute of Technology, funded by the National Aeronautics and Space Administration and the National Science Foundation.

REFERENCES

- Ardila D., Martín E., Basri G., 2000, *AJ*, 120, 479
- Baraffe I., Chabrier G., Allard F., Hauschildt P. H., 1998, *A&A*, 337, 403
- Baraffe I., Chabrier G., Allard F., Hauschildt P. H., 2002, *A&A*, 382, 563
- Barrado y Navascués D., Bouvier J., Stauffer J. R., Lodieu N., McCaughrean M. J., 2002, *A&A*, 395, 813
- Barrado y Navascués D., Martín E. L., 2003, *AJ*, 126, 2997
- Blaauw A., 1964, in Kerr F. J., ed., *IAU Symp. 20: The Galaxy and the Magellanic Clouds The Scorpio-Centaurus association*. p. 50
- Bouy H., Martín E. L., Brandner W., et al. 2006, *A&A*, 451, 177
- Chabrier G., Baraffe I., Allard F., Hauschildt P., 2000, *ApJ*, 542, 464
- de Bruijne J. H. J., Hoogerwerf R., Brown A. G. A., Aguilar L. A., de Zeeuw P. T., 1997, in *ESA SP-402: Hipparcos - Venice '97 Improved Methods for Identifying Moving Groups*. pp 575–578
- de Zeeuw P. T., Hoogerwerf R., de Bruijne J. H. J., Brown A. G. A., Blaauw A., 1999, *AJ*, 117, 354

- Dye S., Warren S. J., Hambly N. C., et al. 2006, *astro-ph/0603608*
- Hamuy M., Walker A. R., Suntzeff N. B., et al. 1992, *PASP*, 104, 533
- Hawley S. L., Gizis J. E., Reid I. N., 1996, *AJ*, 112, 2799
- Hewett P. C., Warren S. J., Leggett S. K., Hodgkin S. T., 2006, *MNRAS*, 367, 454
- Kirkpatrick J. D., Henry T. J., McCarthy D. W., 1991, *ApJS*, 77, 417
- Kirkpatrick J. D., Reid I. N., Liebert J., et al. 1999, *ApJ*, 519, 802
- Lawrence A., Warren S. J., Almaini O., et al. 2006, *astro-ph/0604426*
- Leggett S. K., 1992, *ApJS*, 82, 351
- Luhman K. L., 2005, *ApJL*, 633, L41
- Luhman K. L., Rieke G. H., Lada C. J., Lada E. A., 1998, *ApJ*, 508, 347
- Martín E. L., Delfosse X., Basri G., et al. 1999, *AJ*, 118, 2466
- Martín E. L., Delfosse X., Guieu S., 2004, *AJ*, 127, 449
- Martín E. L., Rebolo R., Zapatero-Osorio M. R., 1996, *ApJ*, 469, 706
- Mohanty S., Jayawardhana R., Basri G., 2005, *ApJ*, 626, 498
- Morau E., Bouvier J., Stauffer J. R., 2001, *A&A*, 367, 211
- Preibisch T., Guenther E., Zinnecker H., et al. 1998, *A&A*, 333, 619
- Preibisch T., Zinnecker H., 2002, *AJ*, 123, 1613
- Prosser C. F., 1992, *AJ*, 103, 488
- Slesnick C. L., Carpenter J. M., Hillenbrand L. A., 2006, *AJ*, 131, 3016
- Walter F. M., Vrba F. J., Mathieu R. D., Brown A., Myers P. C., 1994, *AJ*, 107, 692

APPENDIX A: SQL QUERY SUBMITTED TO THE WFCAM SCIENCE ARCHIVE

Initial sample selection for this Upper Sco study was made by accessing the UKIDSS Early Data Release database (UKIDSSDR) held at the WFCAM Science Archive. The SQL query given in Figure A1 was used.

For more details concerning the use of SQL and data in the WFCAM Science Archive, see Hambly et al. (2006) and references therein.

```

SELECT
/* Attribute selection: */
g.ra, g.dec, zmyPnt, ymjPnt, jmhPnt, hmk_1Pnt,
zaperMag3, yaperMag3, japerMag3, haperMag3, k_1aperMag3,
3.6e6*COS(RADIANS(g.dec))*(g.ra-T2.ra)/((mj.mjdObs - T2.jdate+2400000.5)/365.25) AS pmRA,
3.6e6*(g.dec-T2.dec)/((mj.mjdObs - T2.jdate+2400000.5)/365.25) AS pmDEC

FROM
/* Table(s) from which to select the attributes: */
gcsMergeLog AS l, Multiframe AS mj, (
    SELECT t.ra AS ra, t.dec AS dec, x.slaveObjID AS slaveObjID,
           x.masterObjID AS masterObjID, t.j_m, t.h_m, t.k_m, t.jdate
    FROM   gcsSourceXtwomass_psc AS x, TWOMASS..twomass_psc AS t
    WHERE  x.slaveObjID=t.pts_key AND distanceMins IN (
        SELECT MIN(distanceMins) FROM gcsSourceXtwomass_psc WHERE masterObjID=x.masterObjID
    )
) AS T2 RIGHT OUTER JOIN gcsSource AS g ON (g.sourceID=T2.masterObjID)

WHERE
/* Sample selection predicates:
only Upper Sco data (no other GCS target is south of the equator) */
g.dec < 0.0
/* Bright saturation cut-offs */
AND zaperMag3 > 11.4
AND yaperMag3 > 11.3
AND japerMag3 > 10.5
AND haperMag3 > 10.2
AND k_1aperMag3 > 9.7
/* Limit merged passband selection to +/- 1 arcsec */
AND zXi BETWEEN -1.0 AND +1.0
AND yXi BETWEEN -1.0 AND +1.0
AND jXi BETWEEN -1.0 AND +1.0
AND hXi BETWEEN -1.0 AND +1.0
AND k_1Xi BETWEEN -1.0 AND +1.0
AND zEta BETWEEN -1.0 AND +1.0
AND yEta BETWEEN -1.0 AND +1.0
AND jEta BETWEEN -1.0 AND +1.0
AND hEta BETWEEN -1.0 AND +1.0
AND k_1Eta BETWEEN -1.0 AND +1.0
/* Retain only point-like sources */
AND zClass BETWEEN -2 AND -1
AND yClass BETWEEN -2 AND -1
AND jClass BETWEEN -2 AND -1
AND hClass BETWEEN -2 AND -1
AND k_1Class BETWEEN -2 AND -1
/* Retain only the best record when duplicated in an overlap region */
AND (priOrSec = 0 OR priOrSec = g.frameSetID)
/* Table join predicates: */
AND g.frameSetID=l.frameSetID
AND l.jmfiD=mj.multiframeID

```

Figure A1. Structured Query Language (SQL) query used on the WFCAM Science Archive database UKIDSSDR to select the GCS Upper Sco sample discussed in the paper. The query returns 174,010 rows of data.